

b) h. w. with basal hair pencil (*Loesa Moore*).

The only species of this group lives in Macromalayana and differs from the true *Lohora* in its shorter cell h. w., while the lower discocellular inosculates the furcation of the two median veins, the middle one is also straight.

M. oroatis *Hew.* Type from Java where the species is very rare. The ♀ is still unknown. I have before me four ♂♂ from the West and one from the East of the island and these are identically the same. DOHERTY also found *oroatis* in Bali. *oroatis* is remarkable for a tuft of thin yellowish hair situated just above the centre of submarginal of h. w. This tuft gave rise to the forming of a genus *Loesa*, containing only one species; fully described forms must fall into the *oroatis* group. Java. Dry-season form has not been examined by me. — **ustulata** *Dist.* (93 a) lives at foot of fore-alps in company with *orseis* and *fuscum*; the darkest of the yellow species in Sumatra. The ♂ has a lustrous red-brown, the larger ♀ a dull light brown ground colour on both wings. Malay peninsula, N.E. and W. Sumatra. In Tenasserim appears as a great rarity **surkha** *Marsh.* (= *fervida* *Btlr.*) in two seasonal forms, it is based on the form without eyespots which apparently does not arrive at full development in Sumatra and Java. Moreover the Tenasserim race is so close to *ustulata* that ELWES will only recognize it as a rainless season aberration.