

4. Subgroup.

The hindwing of the ♂ above always with two hair-pencils, one in the cell close to the costal margin and one in the basal part of cellule 6; in the normal position of the wings both are covered by the hindmargin of the forewing. The forewing almost always with a small, raised, weal-like mealy spot in cellule 1a, close behind vein 1. — The upper surface of the wings always without white markings and the forewing never with sharply defined yellow subapical band.

M. golo *Auriv.* (27 e). Both wings above unicolorous dark brown without markings, in the ♀ somewhat lighter than in the ♂: on the under surface the basal part is dark velvety brown without distinct markings and distally almost straight, the median line being only slightly curved on the forewing and nearly straight on the hindwing, this line and the marginal area brown in the type-form; the eye-spots are black, white-pupilled, with a yellow and a black-brown ring; the forewing with a large eye-spot in cellule 2, a small one in 5 and a punctiform in 4 and in 6; the hindwing has 7 eye-spots, of which only the spot in cellule 2 is larger, but the one in cellule 6 not larger than that in cellule 5. ♂: the pencils on the hindwing are light yellowish; the forewing with a small, not raised mealy spot in cellule 1a about at the middle of vein 1. Cameroons, Congo, Uganda. — In **violascens** *Auriv.*, from Gaboon and the Congo, the median line on the under surface is broader and shows, together with the marginal area, a strong violet reflection.