

1103 a. *Eucereon xanthura*. (Plate XIX. fig. 7.)

*Eucereon xanthura*, Schaus, A. M. N. H. (8) vi. p. 192 (1910).

*Eucereon schausi*, Roths. Nov. Zool. xix. p. 175 (1912).

*Eucereon apicaflava*, Kaye, Ent. Rec. xxii. p. 115 (1914).

Head greyish fuscous, the frons with white bar above, the basal joint of antennæ white in front, the back of head with pair of yellow spots; thorax grey mixed with white, the patagia with black streaks and white edges, the dorsum of thorax with pair of white stripes; pectus and legs fuscous, the femora and base of mid tibiæ streaked with white; abdomen fuscous, the extremity yellow, sublateral series of white patches except on terminal segments. Fore wing white, the veins, discal and submedian folds streaked with black; antemedial line with oblique black bar from costa, then slight and angled inwards on median nervure, then almost obsolete and angled outwards at submedian fold; an oblique black lunule from costa to lower angle of cell with some white in centre; postmedial line represented by a series of short black dashes on the veins, oblique from costa to vein 4, then very obliquely incurved to middle of inner margin and with traces of a line between them; subterminal line black, with spot at costa, angled outwards below veins 7 and 6, bent outwards to near termen from above vein 4 to vein 3, then bent inwards to vein 2 and outwards to tornus, at vein 3 connected by an oblique bar with the termen; a black spot on costa before apex. Hind wing whitish tinged with fuscous, the terminal area darker, the veins slightly streaked with black; the underside white with the veins, costal area, and terminal area to vein 1 suffused with fuscous.

*Hab.* MEXICO, Jalapa (*Schaus*), type † ♂ *schausi* in Coll. Rothschild; COSTA RICA, Juan Vinas (*Schaus*), 1 ♀; COLOMBIA, Don Amo (*H. H. Smith*), 3 ♂, Sierra del Libano (*H. H. Smith*), 1 ♀; ECUADOR, Paramba; VENEZUELA, Caracas, type † *apicaflava* in Coll. Kaye; BRAZIL, Rio Janeiro, Tijuca (*Wagner*), 2 ♀. *Exp.* 44 millim.